

Journal of Herbal Medicine

Volume 6, Issue 1, March 2016, Pages 18-23

Research paper

Effect of lavender essential oil as a prophylactic therapy for migraine: A randomized controlled clinical trial

Shahram Rafie a, Forough Namjoyan b, Fereshteh Golfakhrabadi b, Fatemeh Yousefbeyk c, Alireza Hassanzadeh a 💍 🖾

Show more ✓

≪ Share **⋾** Cite

https://doi.org/10.1016/j.hermed.2016.01.003 \nearrow Get rights and content \nearrow

Abstract

There is no cure for migraine, but preventive treatments are usually applied to reduce the frequency and severity of <u>headache</u> attacks. The purpose of this study was to investigate the effect of lavender as a prophylactic therapy for migraine in a randomized <u>controlled clinical trial</u>. This double-blind and placebocontrolled study was conducted over a period of three months. Patients were assessed for migraine impact at the baseline and at the end of the study, using the Migraine Disability Assessment Scores (MIDAS) questionnaire. In the case group, after three months of lavender therapy, the MIDAS score was reduced. The reduction in MIDAS score was significant (P<0.05), when compared to the baseline and also control group. During the treatment, participants did not report any complaints or <u>side effects</u>. The results of this present study report that the frequency and severity of migraine incidents were reduced in those participants using lavender therapy during the three month trial.

Introduction

Migraine is influenced by various factors such as diet, hormonal disorders, digestive problems, autoimmune disturbances, structural imbalances, mental stress and lifestyle (Fantasia, 2014). It is a common and disabling disorder amongst adults and children, although it more often affects women (Magis and Schoenen, 2011). Irrespective of age, migraine can spontaneously occur in an individual, but migraine often starts between ages 10 and 30 (Magis and Schoenen, 2011). The migraine pain is often accompanied by a variety of symptoms which may include nausea, blurred vision, vomiting, sensitivity to light, and noise (Fantasia, 2014).

The following medications are commonly used to prevent migraine: beta-blockers, flunarizine, topiramate, valproate, amitriptyline, venlafaxine, gabapentin, magnesium and botulinum toxin type A (Chayasirisobhon, 2013). None of the drugs used in migraine prophylaxis are uniformly effective for patients. There are minimal but still reported side effects associated with common medications (Chayasirisobhon, 2013, Prior et al., 2010).

There is no cure for migraine, but supplements and medicinal herbs can be used to prevent migraine attacks and reduce the frequency and severity of a headache (D'Amico et al., 2006, Lipton et al., 2003). In addition, the use of traditional medicine and medicinal plants is increasing worldwide. Therefore, the study of their clinical efficacy and the conduct of further clinical trials regarding medicinal herbs is required as a matter of priority. The results of such research can be used for further investigations into pharmaceutical formulations and also to improve our knowledge regarding herbal medicine treatment in its own right. Some medicinal herbs have shown capability in the treatment of migraine prophylaxis. Feverfew (*Tanacetum parthenium*), for example, was found to be effective in the prevention of migraine. The active constituents of feverfew are sesquiterpene and lactones, especially parthenolide (Johnson et al., 1985, Bohlmann and Zdero, 1986, Murphy et al., 1988). Butterbur (*Petasites hybridus* root) has shown efficacy in migraine prophylaxis (Lipton et al., 2004). This plant acts by the inhibition of peptide leukotriene biosynthesis and interference in the inflammatory cascade associated with migraine (Eaton, 1998, Sheftell et al., 2004, Grossman and Schmidrams, 2000).

The genus *Lavandula* (common name: lavender) is comprised of about 25–30 species of flowering plants in the Lamiaceae (Labiatae) family (Behbahani et al., 2013, Effati-Daryani et al., 2015). Native to France and the western Mediterranean these flowering plants are cultivated worldwide for their volatile oil (Behbahani et al., 2013, Effati-Daryani et al., 2015). The leaves of lavender can be used in aromatherapy; lavender aromatherapy shows positive effects on hemodynamic indices among patients with acute coronary syndrome (Nategh et al., 2015). In another study, lavender essential oil was found to reduce the level of anxiety in patients undergoing coronary artery bypass graft surgery (Seifi et al., 2014). A recent evaluation showed the positive effect of lavender essential oil, in reducing anxiety in patients undergoing curettage. It may therefore be concluded that lavender aromatherapy can be used in supportive therapy alongside orthodox treatment (Bakhsha et al., 2014).

Lavender essential oil has been used traditionally for the treatment of colds, digestion, flatulence, upset stomach, liver, gallbladder problems and loss of appetite (Katona et al., 2010, Kim et al., 2007). In addition, this genus is beneficial for stress, anxiety, exhaustion, headaches, migraines, insomnia and depression (Seifi et al., 2014, Katona et al., 2010, Kim et al., 2007). *Lavandula stoechas* L. which is generally called Ustkhuddus in the Persian language, is a low-growing and evergreen herb (Lim, 2014). There are some reports about its spasmolytic (Gedney et al., 2004), sedative (Buchbauer et al., 1991), antihypertensive (Koto et al., 2006), antimicrobial (Inouye et al., 2001) and antifungal (D'Auria et al., 2005) properties. Moreover, the analgesic effect of lavender oil has been investigated in previous studies (Yip and Tse, 2006, Gedney et al., 2004). The efficacy of lavender oil in anxiety disorders has also been reported (Morris, 2002). The purpose of this study was to investigate the effect of lavender essential oil as a prophylactic therapy for migraine in a placebocontrolled clinical trial.

Access through your organization

Check access to the full text by signing in through your organization.

Access through your organization

Section snippets

Preparation of lavender essential oil and placebo

The aerial parts of *L. stoechas* L. were collected from the Zardband Botanical Garden of Iran in May 2014. Identification and authentication was carried out by botanists in Zardband Botanical Garden and a voucher specimen deposited in their herbarium. The samples were crushed and the essential oil was obtained by the hydrodistillation method using a Clevenger type apparatus (Advanced Technocracy Inc., India), according to the European Pharmacopoeia (1975) (Maisonneune, 1975, Golfakhrabadi et ...

Results

In this study, the efficacy of lavender essential oil as a prophylactic therapy for migraine was investigated. Table 1 shows the demographic data of the study population. There were 20 female and 10 male patients in the treatment group and 22 female and 8 male in the control group. The average age of patients in the group receiving lavender was 30.40±8.92 and the average age of patients receiving placebo was 27.70±7.00. There were no significant differences between patients with regard to age ...

Discussion

In this clinical trial, all participants were previously being treated with propranolol and after agreeing to participate in this study, also received lavender extract as an adjunct to that treatment. At the end of the study, the improvement of headache severity and frequency in the case group compared to the control group after adjunct therapy with lavender extract was significant (P<0.05). Headache severity in the case group was reduced to 41.1% in the first month and 52.35% in the third ...

Conclusion

The results of this present study report that the frequency and severity of migraine incidents were reduced in those participants using lavender therapy during the three month trial. Essential oil of lavender shows significant promise for its efficacy in the prevention of migraine however since the use of lavender extract was concomitant with propranolol, the patients' current orthodox treatment, these results must be viewed with caution in regard to possible synergy between the two treatments. ...

Conflict of interest

The authors declare that there is no conflict of interest. ...

Acknowledgement

This research was supported by Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran (Grant number: ajums.REC.1392.214). ...

Recommended articles

References (38)

H.C. Fantasia

Migraine headache prophylaxis in adolescents

Nurs. Women's Health (2014)

J.M. Katona et al.

Microencapsulation of oil by polymer mixture–ionic surfactant interaction induced coacervation Carbohydr. Polym. (2010)

N. Morris

The effects of lavender (*Lavendula angustifolium*) baths on psychological wellbeing: two exploratory randomised control trials

Complement. Ther. Med. (2002)

J.J. Murphy et al.

Randomised double placebo-controlled trial of feverfew in migraine prevention Lancet (1988)

Y.B. Yip et al.

An experimental study on the effectiveness of acupressure with aromatic lavender essential oil for sub-acute, non-specific neck pain in Hong Kong

Complement. Ther. Clin. Pract. (2006)

O. Algoui-Ismaili et al.

Odor hedonics: connection with emotional response estimated by autonomic parameters Chem. Senses (1997)

F. Bakhsha et al.

Phytochemical and anti-oxidant activity of *Lavandula angustifolia* Mill. essential oil on preoperative anxiety in patients undergoing diagnostic curettage

IJWHR (2014)

B.A. Behbahani et al.

Antimicrobial effects of Lavandula stoechas L. and Rosmarinus officinalis L. extracts on Escherichia coli and Staphylococcus aureus

Sci. J. Microbiol. (2013)

F. Bohlmann et al.

Sesquiterpene lactones and other constituents from Tanacetum parthenium

Phytochemistry (1986)

G. Buchbauer et al.

Aromatherapy: evidence for sedative effects of the essential oil of lavender after inhalation

Z. Naturforsch. C (1991)



View more references

Cited by (34)

Efficacy of Anise (Pimpinella anisum L.) oil for migraine headache: A pilot randomized placebocontrolled clinical trial

2019, Journal of Ethnopharmacology

Citation Excerpt:

...However, there are trace evidences for their use in clinical practice. Although, efficacy of several herbal remedies such as lavender essential oil, butterbur, feverfew, ginger, and rose oil were evaluated in previous studies (Diener et al., 2018; Rafie et al., 2016; Martins et al., 2018; Vogler et al., 1998). Of course, most of these herbal drugs have been evaluated in a systemic dosage form....

Show abstract ✓

A new cost-effective approach for lavender essential oils quality assessment

2018, Industrial Crops and Products

Show abstract ∨

Essential oil composition and antioxidant activities of eight cultivars of Lavender and Lavandin from western Anatolia

2018, Industrial Crops and Products

Citation Excerpt:

...Lavandula essential oil, with the main active constituents linalool, linalyl acetate, 1,8-cineole, cis and trans-ocimene, terpinen-4-ol and camphor, has been reported to have antimicrobial, anticholinesterase and antioxidant activities (Costa et al., 2012; Hanamanthagouda et al., 2010; Cavanagh and Wilkinson, 2002; Gonçalves and Romano, 2013). Thus, Lavandula oil promotes healing symptoms for stress, exhaustion, migraines, anxiety, insomnia and depression (Rafie et al., 2016; Danh et al., 2013; Fismer and Pilkington, 2012; Koulivand et al., 2013). When we consider all the pharmacological properties and the rich chemical content, lavandula essential oil is a significant product and encourage the cultivation of this plant as an industrial crop for essential oil production (Stanev et al., 2016; Adal et al., 2015)....

Show abstract ✓

Neuroprotective Potential of Aromatic Herbs: Rosemary, Sage, and Lavender

2022, Frontiers in Neuroscience

Phytochemicals from the genus Lavandula: a review ¬

2021, Phytochemistry Reviews

Essential Oil Compositions and Antibacterial and Antioxidant Activities of Five Lavandula stoechas Cultivars Grown in Thailand

2019, Chemistry and Biodiversity



View all citing articles on Scopus ↗

View full text

Copyright © 2016 Elsevier GmbH. All rights reserved.



All content on this site: Copyright © 2025 or its licensors and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the relevant licensing terms apply.

